





DEPARTMENT OF CON

NOAA

Rick A. Hart, Ph.D.
NOAA Fisheries Service
Southeast Fisheries Science Center
Galveston Laboratory
Galveston, TX USA

Gulf of Mexico Fisheries Management Council April, 2016






Closure

The GMFMC Shrimp FMP implemented in 1981 for the EEZ fishery


- Goal was to increase the yield of brown shrimp harvested from offshore Texas waters
- Historically the closure is from Mid-May to Mid-July
- Since 1990 the near shore < 4 fathoms area has also been closed






Texas brown shrimp offshore catch for July and August, 1981-2015.






Monthly brown shrimp offshore shrimp catch for May – August, 2015.






Landings by upper Texas ports 2015.






Landings by middle Texas ports 2015.






Landings by lower Texas ports 2015.





July offshore white shrimp catch for 2015.





August offshore white shrimp catch for 2015.



Summary

- Environmental factors important for growth and abundance of shrimp – below average this year
- Brown shrimp catch off Texas coast below long-term average
- Brown shrimp size off Texas about 1.5% in the >67 count size category



Summary

- Some changes in shrimp landings distribution in Texas ports
- Increase in pounds yield with 2015 closure between 0 and 7%
- White shrimp catch off Texas below average during both July and August



